

Lignée ORGAPRED N°OV-135_A

Le MIAOU (**M**inimal **I**nformation **A**bout an **O**rganoid and its **U**se):

Eléments descriptifs permettant à l'homme de l'art de reproduire une expérience de fabrication, de caractérisation et d'étude fonctionnelle d'organoïdes

Le MIAOU sert à identifier les informations présentes (la réponse Oui/Non est la plus importante) et à évaluer la qualité de leur description pour la reproductibilité.

A) SOURCE MATERIAL MATERIEL SOURCE

Informed consent obtained Consentement adapté au but de la recherche	Yes
Collection declaration Déclaration de collection- CODECOH*	Yes
Descriptors: gender, age, anatomical region, diagnostic, viral statut Descriptif : genre, âge, région anatomique, diagnostic, statut viral	Yes
Clinical data on the patient Tableau clinique du patient	Yes

* : gestion de la COnservation D'Éléments du COrps Humain

Primary cell of patient (and healthy subjects) and tumors Cellule primaire de patient (et sujets sains) et tumeurs

Genetic identity at arrival (example: DNA sequence, snips, digital PCR, STR, CGH array) Identité génétique à réception	Yes, STR on tumor and organoid sample for quality control
Genetic quality control (example : Karyotype, STR, digital PCR) Contrôle de qualité génétique	Yes, STR on tumor and organoid sample for quality control
Functional quality (example: differentiation test for pluripotency of iPSCs, permeability tests for intestinal epithelial cells...) Qualité fonctionnelle	Yes, organoid establishment
Cell identity after X passages identité cellulaire après X passages	STR analysis on tumor DNA and on organoid DNA at passage 5.
Cell type marker (example : marker name, detection method, target value) Marqueur de type cellulaire	IHC specific markers on tumor fragment: Ki67 / p53 / PAX8 / WT1 / CXCR4 / H3
Number of passages at arrival Nombre de passages à la réception	0, cells directly obtained from patient's tumor
Number of possible or required passages before genesis of organoids Nombre de passages possibles ou requis avant genèse des organoïdes	0, considered established at passage 3 and above
Storage conditions Protocole de conservation	Yes, between 100 and 1000k cells in 2 mL cryovials, resuspended in 1000 µL in freezing solution (10% DMSO, 90% FBS) and

Lignée ORGAPRED N°OV-135_A

	frozen gradually decreasing temperature (1°C/min in CoolCell) to -80°C before long-term storage at -196°C.
Mutations if genetic disease Mutations si maladie génétique	No mutation found by NGS (5 genes panel)
Contamination tests (mycoplasma, bacteriological, fungal) Tests contamination	Absence of bacterial or fungal contamination Absence of Mycoplasma (biochemical test)
Method of tissue dissociation (production of single-cell material or tissue substructures - example: intestinal crypt) Méthode de dissociation du tissu	Yes, 1h enzymatic and mechanical dissociation of tumor fragments with Tumor Dissociation Kit, human (Miltenyi) and 100 µm filtration.

Storage conditions of the lines or cells Conditions de conservation des lignées ou cellules

Master banks, (description of protocols, drift control) Banques mères	Yes, BRC of the BACLESSE Centre, NF S96-900 certified
Daughter banks (description of protocols, drift control) Banques filles....	No
Storage: freezing and thawing protocol Conservation : protocole de congélation et de décongélation	Freezing: between 100 and 1000k cells in 2 mL cryovials, resuspended in 1000 µL in Recovery Cell Culture Freezing Medium (Gibco) and frozen gradually decreasing temperature (1°C/min in CoolCell) to -80°C before long-term storage at -150°C. Thawing: Fast thawing at 37°C, transfer in fresh culture media before centrifugation and matrix seeding.
Storage modalities Modalités de conservation	Long-term storage at -150°C.

B) MANUFACTURING OF THE ORGANOIDS FABRICATION DE L'ORGANOIDE

Culture conditions of cells Conditions de culture des cellules

Composition of culture media, nature, origin and quantities of supplements used (e.g. glucose, serum, antibiotics, growth factors etc.) Composition des milieux	Culture in an enriched medium [Advanced DMEM (Gibco) supplemented with 100 UI/mL of penicillin and streptomycin (Gibco), 1% GlutaMAX (Gibco), 1X B27 (Gibco), 10 mM Nicotinamide (Sigma-Aldrich), 1.25 mM N-Acetyl-L-Cysteine (Sigma-Aldrich), 50 µg/mL Primocin (InvivoGen), 5 µM Y27632 (Interchim), 20 ng/mL FGF-10 (PeproTech), 500 nM A-83-01 (PeproTech), 50 ng/mL EGF (PeproTech), 1 ng/ml FGF-basic (PeproTech), 1 µM SB202190 (PeproTech), 1 µM PGE2 (Sigma-
---	---

Lignée ORGAPRED N°OV-135_A

	Aldrich), 10% RSP01-conditioned media (Cultrex HA-R-Spondin1-Fc 293 T, Amsbio) and 50% L-WRN- conditioned media (Cultrex L-WRN, Amsbio)]
Nature and treatment of the supports Nature et traitement des supports	Yes. 24-well plate Costar: culture treated 6-well plate Nunc: culture treated, Nunclon Delta
Seeding conditions Conditions d'ensemencement	10 000 cells per matrix drop (mix of 70% BME2 and 30% culture media v/v)
Frequency of media changes Fréquence des changements de milieu	Twice a week
CO ₂ / O ₂ Concentration	Yes, ambient O ₂ and 5% CO ₂

Generation of organoids (3D): specificities Génération des organoïdes (3D) : spécificités

Matrix culture Culture en matrice

Nature of the matrix (matrigel, hydrogels, hyaluronic acid, human decellularized matrix etc.) Nature de la matrice	Cultrex Reduced Growth Factor Basement Membrane Extract, Type 2, Pathclear
Matrix concentration Concentration de la matrice	70% matrix in 30% media (v/v)
Preparation method (temperature, polymerization time, drop or layer structure, etc.) Modalité de préparation	50 µL drop of matrix and 30 minutes polymerization at 37°C
Seeding density per matrix volume unit Densité d'ensemencement	10 000 cells per matrix drop
Volume and number of drops of matrix per unit area in the culture medium	50 µL drop per well in 24-well plate, 10 drops of 50 µL per well in 6-well plate
Amount of medium depending on the size of the well Quantité de milieu en fonction de la taille du puits	24-well plate: 500µL 6-well plate : 2000µL
Matrix dissociation method for organoid recovery Méthode de dissociation de la matrice pour la récupération des organoïdes	Mechanical dissociation using cold Advanced DMEM media (Gibco) supplemented with 100 UI/mL of penicillin and streptomycin (Gibco), 1% GlutaMAX (Gibco), 1% BSA (Panreac)
Method of dissociation of organoids for their expansion Méthode de dissociation des organoïdes pour leur expansion	Enzymatic dissociation using TrypLE Express Enzyme (1X) solution, 5-15 minutes incubation at 37°C with gentle agitation every 5 minutes, cold Adv DMEM media (1% BSA, see above) is then add to stop the reaction before centrifugation, numeration and seeding.

Lignée ORGAPRED N°OV-135_A

Culture including multiple cell types *Culture incluant de multiples types cellulaires*

Sequence of co-culturing and adaptation of co-culture media Séquence des mises en co-culture	N.A.
Proportion of cell types Proportion des types cellulaires	N.A.

C) ORGANOID CHARACTERIZATION **CARACTÉRISATION DES ORGANOIDES**

The detailed characterization is project dependent, however some standards emerge

<i>Morphology</i> <i>Structure</i>	
Appearance, size, shape [circularity, tubularity, regularity of contour (budding)] Aspect, taille, forme	Round organoids with defined contours
Opacity/réfringency Opacité/réfringence	Opaque
Intra and inter-organoid homogeneity Homogénéité	Homogeneity in shape and size Heterogeneous expression of some IHC markers
Expected morphological, architectural and ultrastructural features, organization of cell types (identity, proportions, distribution) Particularités morphologiques	N.A.

<i>Molecular Characterisation</i> <i>Caractérisation moléculaire</i>	
Elements of genomics, transcriptomics, metabolomics, proteomics, Eléments de génomique, transcriptomique, métabolomique, protéomique	NGS CGH array 3' RNA seq Genomic Instability Scar (GIScar)
Expected specific molecular markers, epigenetic characteristics Marqueurs moléculaires	IHC specific markers on organoids: Ki67 / p53 / PAX8 / WT1 / CXCR4 / H3

<i>Function</i> <i>Fonction</i>	
Qualitative and (if possible) quantitative functional characteristic Caractéristique fonctionnelle	The RECAP test was performed according to the method described in Thorel et al. (2025) (automated scoring of RAD51 foci in PDOs).
Response to treatments (pharmacological, chemical, physical, hormonal...) the treatment protocol, and evaluation (quantitative or qualitative) of the response are described	Protocol: when PDO reached the size of 75–150 µm in diameter, they will be collected and resuspended in PDO treatment medium (PDO culture medium lacking primocin, Y-27,632 and N-

Lignée ORGAPRED N°OV-135_A

Réponse aux traitements	acetylcysteine) with 2% BME2. 200 PDO per well will be seeded in 100 µL volume in a previously coated (1:1 PDO treatment medium/BME2) white clear bottom 96-well plates (Greiner). Drug solutions will then be prepared in a 2% BME2/PDO treatment medium, added to each well and plates will be transferred to a humidified 37°C/5% CO₂ incubator. During the treatment, PDO will be monitored using IncuCyte S3 ZOOM (Sartorius). One week later, ATP levels will be measured by CellTiter-Glo 3D assay (Promega) and luminescence will be quantified using GloMax Discover Microplate Reader (Promega). The half-maximal inhibitory concentration (IC₅₀) and the area under the dose-response curve (AUC) will be computed for each PDO model. <u>Evaluation of treatment response for:</u> Carboplatin = sensitive Paclitaxel = sensitive Doxorubicine = resistant Gemcitabine = sensitive Olaparib = intermediate Niraparib = sensitive
-------------------------	---

<i>Traceability, organoid drift</i> <i>Traçabilité, dérive des organoïdes</i>	
Traceability of components (batches, suppliers etc.. environments, complements) <i>Traçabilité des composants</i>	Yes, traceability of every components (batches number, expiration dates ...)
Traceability of conditioned media (drift of cells used for conditioning, control of lines as for those at the origin of the organoid), control of at least one of the growth factors) <i>Traçabilité des milieux conditionnés</i>	Traceability of every components (batches number, expiration dates ...) used for conditioned media production. No systematic growth factors quantification in conditioned medium.
Drift criteria (morphological, structural, functional, molecular....) specific to each organoid. Specify indices if applicable <i>Critères de dérive</i>	Morphological, response to treatments, IHC markers, STR
Robustness criterion (same starting cells, same organoid). Specify indices if applicable <i>Critère de robustesse</i>	No

D) USE OF ORGANOIDS UTILISATION DES ORGANOIDES

Lignée ORGAPRED N°OV-135_A

Organoid for basic research *Organoïde en recherche fondamentale*

	No
--	----

Organoid in preclinical research (pharmacology, toxicology, ...) *Organoïde en recherche préclinique (pharmacologie, toxicologie, ...)*

Functional similarity criterion between the organoid and the mimicked organ (battery of controls to be performed with target values) Similarité fonctionnelle	Patient clinical data are available to assess concordance with organoid-based tests. Partial correlation between tumor and organoids markers.
Number of usable passages Applicable for: Preclinical development of a drug candidate (IND file) using organoids Nombre de passages exploitables	Yes, more than 10
Number of usable passages Applicable for: Definition of predictive signatures of responses (companion test) Nombre de passages exploitables	Yes, more than 10
Number of usable passages Applicable for : Validation of a care protocol (specific patient) on a cohort: choice of a therapy Validation d'un Protocole de soin	To be defined

Organoid in clinic (personalized, predictive and regenerative medicine, transplantation) *Organoïde en clinique*

GMP certification, total traceability of the components, qualification of the components for Domain 1: Care protocol (specific patient) (validation of the protocol of use of the organoid for the orientation of the therapeutic choice) - Criterion of similarity between the organoid and the biopsy Certification GMP	No
GMP certification, total traceability of components, qualification of components for Domain 2: Use in regenerative medicine (same as cell and tissue therapies) - Functionality criteria, safety (Derivation of biological material and evaluation of the risk of cancer) Certification GMP	No